


# SIDS/SUIDS Risk Factors





## Not sleeping on back

40%\* of SIDS/SUIDS cases from 2010 to 2015 did not sleep on their backs.

■ Sleeping on their back ■ Not sleeping on their back

## Not having a firm, infant approved surface

100%\* of SIDS/SUIDS cases from 2010 to 2015 did not have a firm, infant approved sleeping surface.




## Loose bedding

80%\* of SIDS/SUIDS cases from 2010 to 2015 had loose bedding in their sleeping area(s).


■ Loose bedding ■ Firm bedding

## Smoke exposure

60%\* of SIDS/SUIDS cases from 2010 to 2015 were exposed to smoke.



■ Exposed to smoke ■ Not exposed to smoke




## Cosleeping

70%\* of SIDS/SUIDS cases from 2010 to 2015 were cosleeping.

■ Cosleeping ■ Not cosleeping

## MOB using drugs or alcohol

50%\* MOB of SIDS/SUID cases from 2010 to 2015 smoked, used drugs or alcohol.



■ Used drugs/alcohol ■ Didn't use drugs/alcohol

\*Data collected based off of ten SIDS/SUIDS cases from 2010 to 2015.

\*\*Data collected from Death Certificates, coroner's reports, autopsies, and medical records.