Yolo County Emergency Medical Services Agency

Protocols

Revised Date: May 1, 2024



# **CRUSH INJURY SYNDROME**

Adult

### Definitions

**Crush Injury**: Injury sustained from a compressive force sufficient to interfere with the normal metabolic function of the involved tissue. May be open or closed.

**Crush Syndrome**: Is a Severe systemic manifestation of trauma and ischemia involving soft tissues, principally skeletal muscle, due to prolonged severe crushing. It leads to increased permeability of the cell membrane and to the release of potassium, enzymes, and myoglobin from within cells. Ischemic renal dysfunction secondary to hypotension and diminished renal perfusion results in acute tubular necrosis and uremia.

**Compartment Syndrome**: Local manifestations of muscle ischemia resulting in compressive forces on a closed space.

# Indication

Patients with full extremity (or more) crushed, pinned, or otherwise immobilized with severely impaired circulation.

### BLS

Assess vital signs  $O_2$ , titrate Sp $O_2$  to  $\geq$  94% SMR if indicated Identify and treat life threatening conditions Control external bleeding Prevent hypothermia Consider early request of air ambulance to the scene

### ALS

Cardiac Monitor, 12-Lead ECG, Waveform EtCO<sub>2</sub>, Vascular Access

\* If possible start treatment prior to release of compression

Fluid Bolus NS 250 mL IV/IO (Prior to release of compression)

• May repeat as needed

For hyperkalemia (peaked T-waves, absent P-waves and/or widened QRS complex)

#### Albuterol 5 mg Nebulized

Run continuously

Calcium Chloride 1gm SIVP/IO over 60 seconds

• Flush tubing with NS prior to administering Sodium Bicarbonate

#### Sodium Bicarbonate 1 mEq/kg IV/IO

Max 100 mEq

# Yolo County Emergency Medical Services Agency

Protocols

Revised Date: May 1, 2024

# Consider

- BLS personnel should wait for ALS personnel before attempting extrication
- Treatment may be compromised by confined space

### **Direction**

- For pediatric patients contact closest Pediatric Trauma Center ED Physician
- Early notification of the receiving hospital is essential for proper triage
- Contact Base Hospital for additional treatment