



# Yolo County Emergency Medical Services Agency

## Protocols

Revised Date: May 1, 2024

<b>NEONATAL RESUSCITATION</b>
<b>BLS</b>
Dry and stimulate Provide warmth Insulate head Place neonate skin to skin with mother if possible Suction nose and mouth with bulb syringe
<b>HR &lt; 100 – Labored breathing and cyanosis</b>
Positive-pressure ventilations 40 - 60 BPM, with room air
<b>HR &lt; 60</b>
CPR Chest compression - ventilation ratio 3:1 Continue until HR > 60
<b>Persistent cyanosis</b>
O <sub>2</sub> via blowby or BVM
<b>ALS</b>
Cardiac Monitor, Waveform EtCO <sub>2</sub> , Vascular Access
<b>HR &lt; 60</b>
<b>Epinephrine (1:10,000) 0.01 mg/kg IV/IO</b> <ul style="list-style-type: none"> <li>• Every 3 - 5 minutes</li> </ul> <b>Fluid Bolus NS 10 mL/kg IV/IO</b> <ul style="list-style-type: none"> <li>• No repeat</li> </ul>
<b>Direction</b>
<ul style="list-style-type: none"> <li>• Transport to closest Receiving Hospital with obstetric services</li> <li>• Transport mother with neonate whenever possible</li> <li>• Contact Base Hospital for additional treatment</li> </ul>