

Yolo County Emergency Medical Services Agency

Quick Reference

Revised Date: June 1, 2023

Medication Profile

Midazolam Hydrochloride

(Versed)

<u>Class</u>:

Short-acting benzodiazepine/CNS agent Sedative-Hypnotic Anticonvulsant

Action:

CNS depressant with muscle relaxant, anticonvulsant, and anterograde amnestic effects. Intensifies activity of gamma-aminobenzoic acid (GABA), a major inhibitory neurotransmitter of the brain, by interfering with its reuptake and promoting its accumulation at neural synapses. Also provides some retrograde amnestic effects, making it useful after cardioversion.

Onset: 1-10 minutes

Peak: 20-60 minutes

Duration: 2-6 hours

Indications:

Contraindications:

Seizures Sedation Hypersensitivity/allergy

Side Effects:

CV: Fluctuations in vital signs, hypotension CNS: Oversedation, headache, retrograde amnesia, euphoria, drowsiness, coma RESP: Respiratory depression, respiratory arrest, cough, laryngospasm GI: Nausea, vomiting, hiccough (diaphragmatic spasm producing a cough/noise) EENT: Blurred vision, diplopia (seeing two objects), nystagmus

Notes:

- Midazolam Hydrochloride is an effective chemical restraint and should be used early in the restraint process especially with patients showing signs of excited delirium.
- Can be given in conjunction with acetaminophen, fentanyl or ketamine.

This document is not a substitute for Protocols and Procedures.